

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
20 March 2003 (20.03.2003)

PCT

(10) International Publication Number  
WO 03/022816 A1(51) International Patent Classification: C07D 209/86,  
C07F 7/00

(21) International Application Number: PCT/US02/25889

(22) International Filing Date: 14 August 2002 (14.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/947,581 6 September 2001 (06.09.2001) US(71) Applicant: EQUISTAR CHEMICALS, LP [US/US];  
One Houston Center, 1221 McKinney Street, Houston, TX  
77010 (US).(72) Inventor: HLATKY, Gregory, G.; 6197 Morrow-Ross-  
burg Road, Morrow, OH 45152 (US).(74) Agents: SCHUCHARDT, Jonathan, L. et al.; Lyondell  
Chemical Company, 3801 West Chester Pike, Newtown  
Square, PA 19073 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GH, GI,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC,  
VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KI, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
European patent (AM, AZ, BY, BG, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

## Published:

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE PREPARATION OF BASE-FREE CARBAZOLID ANIONS

(57) Abstract: Base-free carbazolidones are prepared rapidly and in high yield and purity by reaction of a carbazole reactant with alkyl lithium in the presence of a dibasic ligand to form an intermediate complex, followed by reacting the intermediate complex with alkali metal alkoxide.

WO 03/022816 A1

PROCESS FOR THE PREPARATION OF  
BASE-FREE CARBAZOLIDE ANIONS

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

The present invention pertains to a process for the convenient, high-yield preparation of base-free carbazolidine anions.

BACKGROUND ART

Carbazolidine anions have numerous uses in synthetic organic and organometallic chemistry. As but one example, transition metal complexes of carbazolidines are known to have catalytic activity, i.e. as polymerization catalysts for  $\alpha$ -olefin polymerization as disclosed in U.S. Patent 5,539,124. In the syntheses of these compounds, the metal chloride is allowed to react with two equivalents of carbazolidine anion. However, syntheses of many carbazolidine complexes and carbazole derivatives require the use of substantially base-free carbazolidine anions, i.e. potassium carbazolidine. The preparation of base-free compounds has proven difficult.

For example, the deprotonation of carbazole by butyllithium in a toluene or pentane solvent is slow and incomplete, perhaps due to the possibility of hydrogen bonding in the partially deprotonated reaction mixture. Appler (J. Organometal. Chem., 350 (1988) 217) reported that deprotonation of carbazole by butyllithium in toluene required refluxing at 110°C. With the addition of coordinating solvents such as tetramethylethylenediamine (TMEDA) in tetrahydrofuran (THF) or toluene solvent, deprotonation is more rapid and more complete. However, the product is a base-coordinated complex such as  $[\text{Li}(\text{THF})_2][\text{C}_{12}\text{H}_8\text{N}]$  (Hacker et al., Chem. Ber., 120 (1987) 1533) or  $[\text{Li}(\text{TMEDA})][\text{C}_{12}\text{H}_8\text{N}]$ . In subsequent reaction with Group 4 metal compounds such as  $\text{TiCl}_4$ ,  $\text{ZrCl}_4$ ,  $\text{HfCl}_4$ , the Lewis base remains coordinated in the final product to form  $(\text{C}_{12}\text{H}_8\text{N})_2\text{MCl}_2(\text{L})_2$  compounds where L is THF or  $\frac{1}{2}\text{TMEDA}$ . These products are difficult to characterize and subsequent chemistry is complicated by the presence of the solvating group.

It would be desirable to provide a synthesis of carbazolidine salts which are


free of Lewis bases, producing the carbazolidine salt in reasonable yields and with reasonable reaction times.

### SUMMARY OF THE INVENTION

It has now been surprisingly discovered that carbazolidine ions free of Lewis  
 5 bases and complexing neutral ligands can be prepared rapidly and in high yield by reacting carbazole with a deprotonating agent in hydrocarbon solvent in the presence of at least an equimolar amount of an aprotic coordinating ligand to form an intermediate complex, followed by reaction in hydrocarbon solvent in the presence of one equivalent of alkali alkoxide to yield the base-free alkali  
 10 carbazolidine.

### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT(S)

Any carbazole is believed to be useful in the present process. Preferred carbazoles correspond to the formula



where R<sup>1</sup> through R<sup>8</sup> are hydrogen or substituents which do not interfere with  
 20 obtaining the desired carbazolidine anion product. Preferred R<sup>1</sup> through R<sup>8</sup> are hydrogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, polyoxyalkyl, aryl, heteroaryl, silyl, or two adjacent R<sup>1</sup> through R<sup>8</sup> may form a C<sub>4-6</sub> cycloalkyl or 6-membered aryl or heteroaryl ring or a C<sub>4-6</sub> cycloalkyl or 6 membered aryl or heteroaryl portion of a more extended ring system. Any of R<sup>1</sup> through R<sup>8</sup> and  
 25 more preferably any of R<sup>2</sup> through R<sup>7</sup> and yet more preferably any of R<sup>3</sup> through R<sup>6</sup> may be an alkylene, alkenylene or silyl, etc. bridging linkage between a first carbazolidine anion and a second carbazolidine anion. Any one of C-R<sup>1</sup> to C-R<sup>8</sup> can also be replaced by a nitrogen, phosphorous, or arsenic atom.

Preferably, R<sup>1</sup> through R<sup>8</sup> are individually hydrogen, C<sub>1-4</sub> lower alkyl, or  
 30 one or more pairs of adjacent R<sup>1</sup> through R<sup>8</sup> (including the pair of R<sup>4</sup> and R<sup>5</sup>) may form an aryl or heteroaryl ring or an aryl or a heteroaryl portion of a larger aromatic ring system. Further examples of preferred carbazoles are found in U.S.

5,539,124. When  $R^1$  through  $R^5$  are alkyl, they are preferably  $C_{1-4}$ alkyl, more preferably methyl. It is preferred that  $R^1$  and  $R^5$  are hydrogen.

The hydrocarbon solvent in which the reaction takes place is a non-complexing hydrocarbon solvent which may also contain heteroatoms such as O, S, or N, as long as the heteroatoms do not facilitate strong complexing of the solvent with the carbazolid product. Most preferably, the solvent is an aromatic, aliphatic or cycloaliphatic hydrocarbon solvent. Preferred cycloaliphatic solvents include cyclopentane, cyclohexane, and cycloheptane. Preferred aliphatic solvents include pentane, hexane, heptane and their branched isomers. Preferred aromatic solvents include benzene, toluene, or o-, m-, or p-xylenes. Mixtures of solvents may be used. The preferred solvent is toluene. Coordinating solvents such as diethyl ether, tetrahydrofuran, and similar ether solvents can be used, but the intermediate lithium salt must be isolated from the reaction mixture for use in the second step of the reaction.

The coordinating ligand is a Lewis base-containing ligand, preferably an aprotic nitrogen- or oxygen-containing ligand. Preferred coordinating ligands include monobasic ligands such as tetrahydrofuran, 1,4-dioxane, and pyridine; chelating dibasic ligands such as N,N,N,N-tetramethylethylenediamine and ethylene glycol dimethyl ether; chelating tribasic ligands such as 1,3,5-trioxane and 1,3,5-trimethyl-1,3,5-triazine; and chelating tetrabasic ligands such as 12-crown-4. Dibasic and polybasic ligands containing both nitrogen and oxygen functionalities as well as ligands containing non-nitrogen and -oxygen complexing Lewis base atoms such as P, O, or S, may also serve as coordinating ligands.

The alkali alkoxide is one which is capable of destroying the intermediate lithium carbazolid/dibasic ligand complex. In general, the alkali alkoxide may be selected from sodium, potassium, and cesium alkoxides, but is preferably a potassium alkoxide. The alkoxide ion may have 1 to 10 or more carbon atoms, thus including methoxide, ethoxide, n-propoxide, isopropoxide, n-butoxide, i-butoxide, s-butoxide, t-butoxide, and other ions derived from linear, branched, or cyclic alkanols. The length of the carbon chain should be such that the resultant lithium alkoxide is soluble in aliphatic, cycloaliphatic, or aromatic solvents. Preferably, the alkali alkoxide is potassium t-butoxide.

The reaction may take place in one or more stages and will be described with reference to the preferred preparation using carbazole and TMEDA as the coordinating ligand. First, the carbazole is reacted in hydrocarbon solvent with alkyl lithium, preferably n-butyl lithium, in the presence of an equimolar amount of TMEDA, at a convenient temperature, i.e. from -20° to 50°C, preferably, 0°C to 25°C. The reaction takes place preferably under moisture-free conditions, preferably under an inert gas blanket, i.e. of nitrogen, argon, etc. Solvents should be dried by conventional techniques. Following the reaction, the complex  $[Li(TMEDA)][C_{12}H_8N]$  may be collected as a white crystalline solid, substantially free of starting material. The intermediate may be separated and washed with additional solvent, or may be retained in the reaction solvent without isolation. If isolated, the intermediate is slurried in toluene or other solvent and reacted with one equivalent of potassium t-butoxide. This reaction may be carried out at temperatures from 0°C to reflux, preferably room temperature to reflux. Potassium carbazolate is isolated as a white crystalline solid and optionally washed with additional solvent to remove all traces of lithium alkoxide. Lithium t-butoxide washes away in the solvent, having substantially pure potassium carbazolate. <sup>1</sup>H NMR generally shows no trace of starting material.

Having generally described this invention, a further understanding can be obtained by reference to certain specific examples which are provided herein for purposes of illustration only and are not intended to be limiting unless otherwise specified.

## EXAMPLES

### Deprotonation of Carbazole

#### Example 1:

Carbazole (3.34 g) was dissolved in tetrahydrofuran (50 mL) and butyllithium (8 mL of a 2.5 M solution in hexane) added. After 30 minutes, the solids were filtered off, washed with pentane, and dried. Yield: 4.79 g. This product was suspended in toluene (100 mL) and potassium t-butoxide (1.69 g) added. The mixture was refluxed for two hours and the pure-white solids filtered hot, washed with toluene and pentane and dried. Yield: 3.04 g. The <sup>1</sup>H NMR spectrum of this product showed no residual N-H proton resonances.

Example 2:

- Carbazole (3.34 g) and N,N,N,N-tetramethylethylenediamine (2.32 g) were stirred in toluene (80 mL). Butyllithium (8 mL of a 2.5 M solution in hexane) was added. After 2 hours, the solution was evaporated to 20 mL and pentane added.
- 5 The product was filtered off, washed with pentane and dried. Yield: 5.00 g. This was suspended in toluene (100 mL) and potassium t-butoxide (1.93 g) added. The mixture was refluxed for 2 hours and cooled. The pure-white product was filtered off, washed with toluene and pentane and dried. Yield: 3.35g. The <sup>1</sup>H NMR spectrum of this product showed no residual N-H proton resonances.

10 Comparative Example 3:

- Carbazole (1.67 g) was suspended in toluene (80 mL) and butyllithium (4 mL of a 2.5 M solution in hexane) added. The mixture was vigorously stirred at room temperature for 2 hours, forming a thick brownish gel. The solids were filtered off with difficulty, washed with pentane, and dried. The <sup>1</sup>H NMR spectrum
- 15 of this product showed residual N-H proton resonances at  $\delta$ 9.89.

## Synthesis of Metal Complexes

Example 4:

- The product of Example 1 was slurried in toluene (50 mL) and added to a suspension of zirconium tetrachloride (1.73 g) in toluene (75 mL). The bright-
- 20 yellow suspension was stirred overnight, then refluxed for two hours. The solids were filtered hot, washed with toluene and pentane and dried. Yield: 4.85 g.

Comparative Example 5:

- Carbazole (5.00 g) was suspended in toluene (60 mL) and butyllithium (12 mL of a 2.5 M solution in hexane) added. The mixture was vigorously stirred for
- 25 two hours and additional toluene (60 mL) added, followed by zirconium tetrachloride (3.5 g). The green-brown slurry was stirred overnight, filtered, washed with toluene and pentane, and dried. Yield: 7.53 g.

Comparative Example 6:


- Carbazole (5.01 g) was suspended in toluene (120 mL) and butyllithium (12
- 30 mL of a 2.5 M solution in hexane) added. The mixture was vigorously stirred overnight. To the thick, light-brown slurry was added zirconium tetrachloride (3.5 g) with an additional aliquot of toluene (20 mL) to wash all the ZrCl<sub>4</sub> into the flask. The orange mixture was stirred for six hours, filtered, washed with toluene

and pentane, and dried. Yield: 9.61 g.

Having now fully described the invention, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the invention as set forth herein. The  
5 terms "a" and "an" mean one or more unless indicated otherwise.

## WHAT IS CLAIMED IS

1. A process for the preparation of alkali metal carbazolides substantially free of basic complexing ligands, said process comprising
- 5 a) reacting a carbazole reactant in hydrocarbon solvent with alkyl lithium in the presence of one equivalent or more of an aprotic coordinating ligand, and optionally isolating the intermediate product;
- b) reacting said intermediate product in hydrocarbon solvent with an alkali metal alkoxide, said alkali metal selected from the group consisting
- 10 of sodium, potassium, rubidium, cesium, and mixtures thereof, thereby obtaining the corresponding alkali metal carbazolidide.
2. The process of claim 1, further comprising isolating said alkali metal carbazolidide as a solid product.
3. The process of claim 2 further comprising washing said solid
- 15 product with non-complexing organic solvent(s) to obtain a purified alkali metal carbazolidide.
4. The process of claim 1, wherein said carbazole has the formula



- where  $R^1$  through  $R^8$  are individually hydrogen, alkyl groups, or other substituents
- 25 which do not interfere with deprotonation of said carbazole with alkyl lithium, or where any two adjacent pairs of  $R^1$  through  $R^8$  together comprise a five or six membered ring system, optionally containing one or more heteroatoms, said ring system optionally being aromatic.
5. The process of claim 4, wherein each of  $R^1$  through  $R^8$  is
- 30 individually selected from hydrogen,  $C_{1-4}$  lower alkyl, silyl,  $C_{1-4}$  alkoxy, cyano, nitro, or where one or more adjacent pairs of  $R^1$  through  $R^8$  together comprise a  $C_{5-6}$  saturated or unsaturated cycloaliphatic ring, or a  $C_3$  or  $C_6$  aryl or heteroaryl



ring, said cycloaliphatic ring(s), aryl ring(s), and heteroaryl ring(s) optionally substituted with non-interfering substituents, and optionally forming part of a larger alicyclic, aromatic, or alicyclicaromatic ring system.

6. The process of claim 1 wherein said alkyl lithium comprises n-butyl  
5 lithium.

7. The process of claim 1 wherein said alkali alkoxide comprises a potassium alkoxide.

8. The process of claim 1, wherein said alkali alkoxide comprises potassium t-butoxide.

10 9. The process of claim 1 wherein said non-complexing organic solvent is a hydrocarbon solvent.

10. The process of claim 1 wherein said non-complexing solvent comprises toluene, xylene, or mixtures thereof.

11. The process of claim 1, wherein said carbazole reactant comprises  
15 carbazole, said alkyl lithium comprises n-butyl lithium, said alkali alkoxide comprises potassium t-butoxide, and wherein said solvent is a liquid aromatic hydrocarbon.

## INTERNATIONAL SEARCH REPORT

 Internal application No  
 PCT/US 02/25889

 A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 7 C07D209/86 C07F7/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHRISTOPH LAMBERT ET AL.: "A novel six-center deprotonation-lithiation reaction mechanism supported by the X-ray structure analysis of a lithium carbazolidine" ANGEWANDTE CHEMIE. INTERNATIONAL EDITION., vol. 31, no. 9, - 1992 pages 1209-1210, XP002229395 VERLAG CHEMIE. WEINHEIM., DE ISSN: 0570-0833 page 1209	1
A	US 5 539 124 A (BRADLEY P. ETHEBERTON ET AL.) 23 July 1996 (1996-07-23) cited in the application column 2 -column 5; example 16	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

30 January 2003

Date of mailing of the international search report

14/02/2003

 Name and mailing address of the ISA  
 European Patent Office, P.B. 5518 Patentkanen 2  
 NL - 2200 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

# INTERNATIONAL SEARCH REPORT

In on on patent family members

Internat pplication No

PCT/US 02/25889

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5539124	A	23-07-1996	
		AU 4609896 A	19-07-1996
		BR 9510288 A	15-12-1998
		CA 2208100 A1	04-07-1996
		CN 1173186 A	11-02-1998
		DE 69520450 D1	26-04-2001
		DE 69520450 T2	06-09-2001
		EP 0799250 A2	08-10-1997
		ES 2157355 T3	16-08-2001
		FI 973660 A	11-09-1997
		JP 11507084 T	22-06-1999
		RU 2164228 C2	20-03-2001
		TW 413685 B	01-12-2000
		WO 9620223 A2	04-07-1996